

EXACTLY FOURTEEN INTRINSICALLY KNOTTED GRAPHS HAVE 21 EDGES

MIN JUNG LEE, HYOUNG JUN KIM, HWA JEONG LEE, AND SEUNGSANG OH

ABSTRACT. Johnson, Kidwell, and Michael showed that intrinsically knotted graphs have at least 21 edges. Also it is known that K_7 and the thirteen graphs obtained from K_7 by ∇Y moves are intrinsically knotted graphs with 21 edges. In this paper, we proved that only these 14 graphs are intrinsically knotted graph with 21 edges.

1. INTRODUCTION

Throughout the article we will take an embedded graph to mean a graph embedded in R^3 . We call a graph G *intrinsically knotted* if every embedding of the graph contains a knotted cycle. Conway-Gordon [2] showed that K_7 , the complete graph with seven vertices, is an intrinsically knotted graph. A graph H is *minor* of another graph G if it can be obtained from G by contracting or deleting some edges. Provided that a graph G is intrinsically knotted and has no proper minor which is intrinsically knotted, G is said to be *minor minimal intrinsically knotted*. Robertson and Seymour [11] proved that there are only finite minor minimal intrinsically knotted graphs, but finding the complete set of them is still an open problem. However, it is well known that K_7 and the thirteen graphs obtained from this graph by ∇Y moves are minor minimal intrinsically knotted [2, 8].

A ∇Y move is an exchanging operation on a graph that is removing all edges of a triangle abc and then adding a new vertex v and three new edges va, vb and vc as in Figure 1. Its reverse operation is called a $Y\nabla$ move. Since ∇Y move preserves intrinsical knottedness [9], we will only consider triangle-free graphs in the article.




FIGURE 1. ∇Y and $Y\nabla$ moves

From the work of Johnson, Kidwell, and Michael [7] it follows that any intrinsically knotted graph consists at least 21 edges. The following two theorems give us the complete set of minor minimal intrinsically knotted graphs with 21 edges.

Theorem 1. *The only triangle-free intrinsically knotted graphs with 21 edges are H_{12} and C_{14} . (H_{12} and C_{14} was described by Kohara and Suzuki in [8].)*

2010 Mathematics Subject Classification: 57M25, 57M27.

This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology (No. 2011-0021795).

The third author was supported by the National Research Foundation of Korea Grant funded by the Korean Government (NRF-2010-0024630).

Kohara and Suzuki [8] found fourteen intrinsically knotted graphs. Goldberg, Mattman, and Naimi [5] constructed twenty graphs derived from H_{12} and C_{14} by $Y\nabla$ moves as in Figure 2, and they showed that these six graphs N_9 , N_{10} , N_{11} , N'_{10} , N'_{11} , and N'_{12} are not intrinsically knotted. This fact was proved by Hanaki, Nikkuni, Taniyama, and Yamazaki [6] independently.

Theorem 2. *Only K_7 and the thirteen graphs obtained from K_7 by ∇Y moves are intrinsically knotted graphs with 21 edges.*




FIGURE 2. K_7 and 19 more relative graphs

2. TERMINOLOGY

From now on let $G = (V, E)$ denote a triangle-free graph with 21 edges. Here V and E denote the sets of all vertices and edges of G , respectively. For any two

distinct vertices a and b , let $\widehat{G}_{a,b} = (\widehat{V}_{a,b}, \widehat{E}_{a,b})$ denote the graph obtained from G by deleting two vertices a and b , and then contracting one edge incident to a vertex of degree 1 or 2 repeatedly until no vertices of degree 1 or 2 exist. Removing vertices means deleting interiors of all edges incident to these vertices and resulting isolated vertices.

In a graph, the distance between two vertices a and b says the number of edges in the shortest path connecting them and is denoted by $\text{dist}(a, b)$. The degree of a vertex a is denoted by $\deg(a)$. To count the number of edges of $\widehat{G}_{a,b}$, we have some notations.

- $E(a)$ is the set of edges which are incident to a .
- $V(a) = \{c \in V \mid \text{dist}(a, c) = 1\}$
- $V_n(a) = \{c \in V \mid \text{dist}(a, c) = n, \deg(c) = n, \}$
- $V_n(a, b) = V_n(a) \cap V_n(b)$
- $V_Y(a, b) = \{c \in V \mid \exists d \in V_3(a, b) \text{ such that } V_3(d) = \{a, b, c\}\}$

Obviously in $G \setminus \{a, b\}$ for some distinct vertices a and b , each vertex of $V_3(a, b)$ has degree 1 and each vertex of $V_3(a), V_3(b)$ (but $V_3(a, b)$) and $V_4(a, b)$ has degree 2. Therefore to derive $\widehat{G}_{a,b}$ all edges incident to a, b and $V_3(a, b)$ are deleted from G , followed by contracting one of the remaining two edges incident to each vertex of $V_3(a), V_3(b), V_4(a, b)$ and $V_Y(a, b)$ as dotted lines in Figure 3(a). Thus we have the following equation counting the number of edges of $\widehat{G}_{a,b}$ which is called a *count equation*;

$$|\widehat{E}_{a,b}| = 21 - |E(a) \cup E(b)| - (|V_3(a)| + |V_3(b)| - |V_3(a, b)| + |V_4(a, b)| + |V_Y(a, b)|)$$

For short, $NE(a, b) = |E(a) \cup E(b)|$ and $NV_3(a, b) = |V_3(a)| + |V_3(b)| - |V_3(a, b)|$. Provided that a and b are adjacent vertices (i.e. $\text{dist}(a, b) = 1$), all of $V_3(a, b), V_4(a, b)$ and $V_Y(a, b)$ are empty because G is triangle-free. Note that this manner of deriving $\widehat{G}_{a,b}$ must be handled in slightly different way when there is a vertex c in V such that more than one vertices of $V(c)$ are contained in $V_3(a, b)$ as in Figure 3(b). In this case we usually delete or contract more edges even though c is not in $V_Y(a, b)$.




FIGURE 3. Deriving $\widehat{G}_{a,b}$

A graph is n -apex if one can remove n vertices from the graph to obtain a planar graph. The following lemma gives an important condition for a graph not to be intrinsically knotted.

Lemma 3. [1, 10] *If G is a 2-apex, then G is not intrinsically knotted.*

The following two lemmas play an important role for G to be 2-apex.

Lemma 4. *If $|\widehat{E}_{a,b}| \leq 8$, then $\widehat{G}_{a,b}$ is a planar graph. Thus G is not intrinsically knotted.*

Lemma 5. *If $|\widehat{E}_{a,b}| = 9$, then $\widehat{G}_{a,b}$ is either a planar graph or homeomorphic to $K(3,3)$. Furthermore if $\widehat{G}_{a,b}$ is not homeomorphic to $K(3,3)$, then G is not intrinsically knotted.*

Note that $K(3,3)$ is a triangle-free graph and every vertex has degree 3.

To prove Theorem 1, we will show that any triangle-free graph with 21 edges is eventually either a 2-apex or homeomorphic to one of H_{12} and C_{14} . Since intrinsically knotted graphs have at least 21 edges [7], it is sufficient to consider simple and connected graphs having no vertex of degree 1 or 2. Our process is constructing all possible such triangle-free graph G with 21 edges, deleting two suitable vertices a and b of G , and then counting the number of edges of $\widehat{G}_{a,b}$. If $\widehat{G}_{a,b}$ has 9 edges or less, we could use Lemma 4 or 5 in order to show that G is not intrinsically knotted. In the event that $\widehat{G}_{a,b}$ is not planar, we will show that G is homeomorphic to H_{12} or C_{14} .

Before describing the proof of Theorem 1, we introduce more notations. Since G is a triangle-free graph with 21 edges, for any vertex a of G , no two vertices in $V(a)$ are adjacent. This means that $E(b)$ and $E(c)$ does not contain an edge in common for any two distinct vertices b and c in $V(a)$.

- $E^2(a) = \bigcup_{b \in V(a)} E(b)$
- $E \setminus E^2(a) = \{e_1(a), \dots, e_{21-n}(a)\}$ if $|E^2(a)| = n < 21$

Sometimes we call $e_i(a)$ an *extra edge* without referring a and two endpoints of the edge are denoted as $x_i(a)$ and $y_i(a)$ where $\deg(x_i(a)) \geq \deg(y_i(a))$.

For visualizing G , we basically proceed the following steps. First choose a vertex a with maximal degree among all vertices and draw $E^2(a)$. If $|E^2(a)| < 21$, draw $E \setminus E^2(a)$ apart from $E^2(a)$ as in Figure 4(a). Then all vertices of degree 1 of $E^2(a)$ and $E \setminus E^2(a)$ are merged into some vertices of degree at least 3 without adding new edges as in Figure 4(b). Let $\overline{V}(a)$ denote the set of all such vertices and $[\overline{V}(a)]$ denote a sequence of the degree of vertices in $\overline{V}(a)$ as follows;

- $\overline{V}(a) = V \setminus (V(a) \cup \{a\}) = \{\overline{v}_1(a), \dots, \overline{v}_m(a)\}$ for $\deg(\overline{v}_i(a)) \geq \deg(\overline{v}_{i+1}(a))$
- $[\overline{V}(a)] = [\deg(\overline{v}_1(a)), \dots, \deg(\overline{v}_m(a))]$
- $|\overline{V}(a)| = \deg(\overline{v}_1(a)) + \dots + \deg(\overline{v}_m(a))$

The graph in Figure 4(b) is an example satisfying $\deg(a) = 5$, $|V_3(a)| = 1$, $|E^2(a)| = 19$, and $[\overline{V}(a)] = [4, 4, 4, 3, 3]$.




FIGURE 4. Visualization of G

The remaining three sections of the article is devoted to proof of Theorem 1. From now on a denotes one of vertices with maximal degree in G . The proof is divided into three parts according to the degree of a . In Section 3 we show that any

graph G with $\deg(a) \geq 5$ cannot be intrinsically knotted. In Section 4 we show that an intrinsically knotted graph with $\deg(a) = 4$ is exactly H_{12} . Finally, in Section 5 we show that any intrinsically knotted graph all of whose vertices have degree 3 is always C_{14} .

3. $\deg(a) \geq 5$

In this section we will show that for some $a', b' \in V$ either $|\widehat{E}_{a', b'}| \leq 8$ or $|\widehat{E}_{a', b'}| = 9$ but $\widehat{G}_{a', b'}$ is not homeomorphic to $K(3, 3)$ by showing that it contains a vertex of degree more than 3 or a triangle (or sometimes a bigon). Then as a conclusion G is not intrinsically knotted by Lemma 4 and 5. Recall that G has 21 edges, every vertex has degree at least 3 and a has the maximal degree among them.

3.1. Case $\deg(a) \geq 6$ or $\deg(a) = 5$ with $|V_3(a)| \geq 4$.

If $\deg(a) \geq 6$ then $|V_3(a)| \geq 3$. Let c be any vertex in $V_3(a)$. Choose a vertex b which has the maximal degree among $V(c) \setminus \{a\}$. Then $|E(b)| + |V_Y(a, b)| \geq 4$ since $|V_Y(a, b)| \geq 1$ when $\deg(b) = 3$. Note that $|V_3(b)| \geq |V_3(a, b)|$. By the count equation, $|\widehat{E}_{a, b}| \leq 8$ in $\widehat{G}_{a, b}$.

Suppose that $\deg(a) = 5$ and $|V_3(a)| \geq 4$. The proof is similar to the previous paragraph.

3.2. Case $\deg(a) = 5$ and $|V_3(a)| = 3$.

Let b and c be two vertices of $V(a) \setminus V_3(a)$. First suppose that both of them have degree 5. Then $NE(a, b) = 9$ and $|V_3(a)| = 3$, so $|\widehat{E}_{a, b}| \leq 9$. Furthermore $\widehat{G}_{a, b}$ has the vertex c of degree 4, so it is not homeomorphic to $K(3, 3)$. Thus G is not intrinsically knotted by Lemma 5.

Now assume that one of them, say b , has degree 4. If $V(b) \setminus \{a\}$ consists of three vertices all of which are of degree 3, then $NE(a, b) = 8$ and $NV_3(a, b) = 6$, so $|\widehat{E}_{a, b}| \leq 7$. If not, let d be a vertex of $V(b)$ which has degree at least 4. Then $NE(a, d) \geq 9$, $|V_3(a)| = 3$, and $|V_4(a, d)| \geq 1$ because $V_4(a, d) \ni b$. This implies that $|\widehat{E}_{a, d}| \leq 8$.

3.3. Case $\deg(a) = 5$ and $|V_3(a)| = 0$.

First suppose that $V(a)$ contains a vertex of degree 5, say c . Since G has 21 edges, all of the other four vertices of $V(a)$ have degree 4. By the previous cases, it is sufficient to consider that $|V_3(c)| \leq 2$. So $V(c) \setminus \{a\}$ has at least two vertices, say b and d , of degree 4 or 5. Since $|E^2(a)| = 21$ and G is triangle-free, all edges of $E(b)$ must be incident to different vertices of $V(a)$, so $|V_4(a, b)| \geq 3$. This implies that $|\widehat{E}_{a, b}| \leq 9$. Since $\widehat{G}_{a, b}$ has the vertex d of degree at least 4, it is not homeomorphic to $K(3, 3)$.

Now assume that all vertices of $V(a)$ have degree 4 providing $|E^2(a)| = 20$. Let $e_1(a)$ be the extra edge and recall that two endpoints of $e_1(a)$ are $x_1(a)$ and $y_1(a)$ with $\deg(x_1(a)) \geq \deg(y_1(a))$. Since G is triangle-free, all edges of $E(x_1(a)) \cup E(y_1(a))$ except $e_1(a)$ must be incident to different vertices of $V(a)$. Thus the degrees of $x_1(a)$ and $y_1(a)$ must be either 4 and 3 or 3 and 3 respectively. If $\deg(x_1(a)) = 4$, then $|V_4(a, x_1(a))| = 3$ and $|V_3(x_1(a))| = 1$, so $|\widehat{E}_{a, x_1(a)}| = 8$. If not, $[\overline{V}(a)]$ is either $[5, 3, 3, 3, 3]$ or $[4, 4, 3, 3, 3]$ because $|\overline{V}(a)| = 17$. Thus $\overline{v}_1(a)$ has degree 5 or 4 and differs from $x_1(a)$ and $y_1(a)$, so $|V_4(a, \overline{v}_1(a))| \geq 4$. Therefore $|\widehat{E}_{a, \overline{v}_1(a)}| \leq 8$.

3.4. Case $\deg(a) = 5$ and $|V_3(a)| = 1$.

In this case $V(a)$ contains four vertices of degree 4 or 5. Let n be the number of such vertices of degree 4, and so $4 - n$ of degree 5 where $n = 2, 3, 4$. This provides that $|E^2(a)| = 21 + (2 - n)$ and $n - 2$ extra edges exist. If $\bar{V}(a)$ contains a vertex $\bar{v}_1(a)$ of degree 5, then five edges of $E(\bar{v}_1(a))$ are extra edges or incident to different vertices in $V(a)$. In any case of n , at least two among these edges are incident to vertices of degree 4 in $V(a)$. Then $NE(a, \bar{v}_1(a)) = 10$, $|V_3(a)| = 1$, and $|V_4(a, \bar{v}_1(a))| \geq 2$, providing $|\hat{E}_{a, \bar{v}_1(a)}| \leq 8$.

Now suppose that $\bar{V}(a)$ contains vertices of degree 3 or 4 only. If $n = 2$, $|\bar{V}(a)| = 16$, and so $[\bar{V}(a)]$ is either $[4, 4, 4, 4]$ or $[4, 3, 3, 3, 3]$. For any vertex b in $V_5(a)$, four edges of $E(b)$ must be incident to different vertices of $\bar{V}(a)$. Indeed these four edges are incident to four vertices of degree 4 or at least three edges among them are incident to vertices of degree 3 in $\bar{V}(a)$. This means that the vertex b has degree 5 with either $V_3(b) = 0$ or $V_3(b) \geq 3$. Both cases are dealt in previous cases (3.3), (3.1), and (3.2).

If $n = 3$, $|\bar{V}(a)| = 17$, and so $[\bar{V}(a)] = [4, 4, 3, 3, 3]$. Let $V_5(a) = \{b\}$. To avoid the case (3.2), four edges of $E(b)$ must be incident to two vertices of degree 4 and two vertices of degree 3 in $\bar{V}(a)$ which are $\bar{v}_1(a)$, $\bar{v}_2(a)$, $\bar{v}_3(a)$, and $\bar{v}_4(a)$. Then there is a vertex c of $V_4(a)$ such that at most one edge of $E(c)$ is incident to $\bar{v}_3(a)$ and $\bar{v}_4(a)$, i.e. two edges of $E(c)$ are incident to $\bar{v}_1(a)$, $\bar{v}_2(a)$, or $\bar{v}_5(a)$. This implies that $NE(b, c) = 9$ and $NV_3(b, c) + |V_4(b, c)| \geq 4$, providing $|\hat{E}_{b, c}| \leq 8$.


Finally if $n = 4$, $|\bar{V}(a)| = 18$, and so $[\bar{V}(a)]$ is either $[4, 4, 4, 3, 3]$ or $[3, 3, 3, 3, 3, 3]$. Recall that two extra edges exist. In the former case let $\{\bar{v}_1(a), \bar{v}_2(a), \bar{v}_3(a)\}$ be the three vertices of degree 4 in $\bar{V}(a)$. For each $i = 1, 2, 3$, if more than two edges of $E(\bar{v}_i(a))$ are incident to $V_4(a)$, then $NE(a, \bar{v}_i(a)) = 9$, $|V_3(a)| = 1$, and $|V_4(a, \bar{v}_i(a))| \geq 3$, providing $|\hat{E}_{a, \bar{v}_i(a)}| \leq 8$. So each of at least two edges of $E(\bar{v}_i(a))$ must be either incident to the unique vertex of $V_3(a)$ or an extra edge. Since G is triangle-free, one of three vertices, say $\bar{v}_1(a)$, has the property that $E(\bar{v}_1(a))$ contains both extra edges, and $V(\bar{v}_1(a))$ and $V(\bar{v}_i(a))$ for each $i = 2, 3$ cannot share a vertex in $V(a)$. This implies that $V(\bar{v}_2(a))$ and $V(\bar{v}_3(a))$ are coincide as in Figure 5(a). Then $NE(\bar{v}_2(a), \bar{v}_3(a)) = 8$, and either $|V_4(\bar{v}_2(a), \bar{v}_3(a))| = 4$ or $|V_4(\bar{v}_2(a), \bar{v}_3(a))| = 3$ and $|V_3(\bar{v}_2(a))| = 1$. Thus $|\hat{E}_{\bar{v}_2(a), \bar{v}_3(a)}| \leq 9$. In $\hat{G}_{\bar{v}_2(a), \bar{v}_3(a)}$ the vertex a still has degree 4 or 5 so that it is not homeomorphic to $K(3, 3)$.

In the latter case let $V_4(a) = \{b_1, b_2, b_3, b_4\}$. We claim that for some $i, j = 1, 2, 3, 4$, $|V_3(b_i, b_j)| \leq 1$. Suppose not, $|V_3(b_i, b_j)| \geq 2$ for all combinations of i and j . By some combinatorics we can derive that all 12 edges of $E(b_1) \cup E(b_2) \cup E(b_3) \cup E(b_4) \setminus E(a)$ are incident to only four vertices of $\bar{V}(a)$ as in Figure 5(b). This means that two extra edges must be incident to the remaining two vertices of $\bar{V}(a)$ at both endpoints. But a bigon is not allowed. Therefore without loss of generality $|V_3(b_1, b_2)| \leq 1$. Then $NE(b_1, b_2) = 8$ and $NV_3(b_1, b_2) \geq 5$, providing $|\hat{E}_{b_1, b_2}| \leq 8$.

3.5. Case $\deg(a) = 5$ and $|V_3(a)| = 2$.

If $V(a)$ contains a vertex of degree 5, say b , then previous four cases guarantee that we only consider that $|V_3(b)| = 2$, so $NV_3(a, b) = 4$ which provides $|\hat{E}_{a, b}| = 8$. Therefore we assume that $V(a)$ contains three vertices of degree 4. In this case three extra edges exist. Since $|\bar{V}(a)| = 19$, $[\bar{V}(a)]$ is one of $[5, 5, 5, 4]$, $[5, 5, 3, 3, 3]$, $[5, 4, 4, 3, 3]$, $[4, 4, 4, 4, 3]$, or $[4, 3, 3, 3, 3, 3]$.

If for some vertex $\bar{v}_i(a)$ with degree 5 one edge of $E(\bar{v}_i(a))$ is incident to $V_4(a)$, then $NE(a, \bar{v}_i(a)) = 10$, $|V_3(a)| = 2$, and $|V_4(a, \bar{v}_i(a))| \geq 1$, providing $|\hat{E}_{a, \bar{v}_i(a)}| \leq 8$. Thus three edges of $E(\bar{v}_i(a))$ are extra edges and the remaining two edges are incident to $V_3(a)$. In the first two cases $[5, 5, 5, 4]$ and $[5, 5, 3, 3, 3]$, both $E(\bar{v}_1(a))$

FIGURE 5. $[4, 4, 4, 3, 3]$ and $[3, 3, 3, 3, 3, 3]$ cases

and $E(\bar{v}_2(a))$ share three extra edges, but G does not have a bigon. In the third case $[5, 4, 4, 3, 3]$, $E(\bar{v}_1(a))$ contains three extra edges and one of these extra edges must be incident to $\bar{v}_4(a)$ or $\bar{v}_5(a)$ both of which have degree 3. Then $NE(a, \bar{v}_1(a)) = 10$ and $NV_3(a, \bar{v}_1(a)) \geq 3$, providing $|\hat{E}_{a, \bar{v}_1(a)}| \leq 8$.

If for some vertex $\bar{v}_i(a)$ with degree 4 two edges of $E(\bar{v}_i(a))$ are incident to $V_4(a)$, then $NE(a, \bar{v}_i(a)) = 9$, $|V_3(a)| = 2$, and $|V_4(a, \bar{v}_i(a))| \geq 2$, providing $|\hat{E}_{a, \bar{v}_i(a)}| \leq 8$. Thus at most one edge of $E(\bar{v}_i(a))$ is incident to $V_4(a)$. In the case $[4, 4, 4, 4, 3]$, at least twelve among sixteen edges incident to four vertices of degree 4 in $\bar{V}(a)$ are not incident to $V_4(a)$. It is impossible because there are only two vertices in $V_3(a)$ and three extra edges. In the last case $[4, 3, 3, 3, 3, 3]$, since only one edge of $E(\bar{v}_1(a))$ is possibly incident to $V_4(a)$, there is a vertex b in $V_4(a)$ such that three edges of $E(b)$ are incident to vertices of degree 3 in $\bar{V}(a)$. Then $NE(a, b) = 8$ and $NV_3(a, b) \geq 5$, providing $|\hat{E}_{a, b}| \leq 8$.

4. $\deg(a) = 4$

Since $|V| = |V_4| + |V_3|$ and $4|V_4| + 3|V_3| = 2|E|$, the pair $(|V_4|, |V_3|)$ has three choices $(3, 10)$, $(6, 6)$ and $(9, 2)$. Here V_n denotes the set of vertices of degree n . As in the preceding section we will show that for some $a', b' \in V$ either $|\hat{E}_{a', b'}| \leq 8$ or $|\hat{E}_{a', b'}| = 9$ but $\hat{G}_{a', b'}$ is not homeomorphic to $K(3, 3)$, providing that G is not intrinsically knotted. But one exception occurs so that G possibly be H_{12} when $(|V_4|, |V_3|) = (6, 6)$.

4.1. Case $(|V_4|, |V_3|) = (3, 10)$.

First suppose that V_4 has a vertex a such that all four vertices of $V(a)$ have degree 3. Let b_1 and b_2 be the other vertices of V_4 . For each $i = 1, 2$, $NE(a, b_i) = 8$. If there is a vertex of $V_3(b_i)$ which is not contained in $V(a)$, then $NV_3(a, b_i) \geq 5$, providing $|\hat{E}_{a, b_i}| \leq 8$. Thus each vertex of $V(b_1)$ is the vertex b_2 or contained in $V(a)$, and similarly for b_2 . This implies that the number of vertices of V_3 which have distance 1 or 2 from the vertex a is at most 6. Take a vertex c of V_3 with distance at least 3 from a . Since each vertex of $V(c)$ is neither b_1 nor b_2 , it has degree 3. Thus $NE(a, c) = 7$ and $NV_3(a, c) \geq 7$, providing $|\hat{E}_{a, c}| \leq 7$.

Suppose not, then without loss of generality three vertices a , b , and c of V_4 have the situation that $V(b)$ contains a and c . If $V_3(a)$ and $V_3(c)$ are not coincide, then $|V_4(a, c)| = 1$ and $NV_3(a, c) \geq 4$, providing $|\hat{E}_{a, c}| \leq 8$. If $V_3(a)$ and $V_3(c)$ are coincide and $|V_Y(a, c)| \geq 2$, then $|V_4(a, c)| = 1$ and $NV_3(a, c) = 3$, providing $|\hat{E}_{a, c}| \leq 7$. If not, for the unique vertex d of $V_Y(a, c)$, $V_3(a) = V_3(c) = V(d)$.

Then for a vertex b' of $V_3(b)$, $V_3(b')$ is disjoint from $V_3(a)$. Thus $NE(a, b') = 7$, $NV_3(a, b') = 5$, and $|V_4(a, b')| = 1$, providing $|\widehat{E}_{a, b'}| \leq 8$.

4.2. Case $(|V_4|, |V_3|) = (6, 6)$.

Consider the subgraph H of G consisting of all edges whose both end vertices have degree 4. Since G has six vertices of degree 3 and the same number of vertices of degree 4, H is not empty set.

Claim 1. If H has a vertex of degree 1, then G is not intrinsically knotted.

Proof. Suppose that H has a vertex a of degree 1. Let b be the unique vertex of degree 4 in $V(a)$. If $|V_3(b)| = 3$, then $NE(a, b) = 7$ and $NV_3(a, b) = 6$, providing $|\widehat{E}_{a, b}| \leq 8$. Thus there is another vertex c of $V_4(b)$ and let $V(c) = \{b, d_1, d_2, d_3\}$. First, assume that $|V_3(c)| = 0$. So the two vertices of $V(b) \setminus \{a, c\}$ must have degree 3 because six vertices a, b, c, d_1, d_2 , and d_3 in V_4 are all different. Thus $NE(a, b) = 7$ and $NV_3(a, b) = 5$, so $|\widehat{E}_{a, b}| \leq 9$. Since $\widehat{G}_{a, b}$ has another vertex d_1 of degree 4, it is not homeomorphic to $K(3, 3)$.

Second, assume that $|V_3(c)| = 1$, say $d_1 \in V_3(c)$. If d_1 is not one of $V(a)$, then $NE(a, c) = 8$ and $NV_3(a, c) + |V_4(a, c)| = 5$, providing $|\widehat{E}_{a, c}| \leq 8$. So we may assume that d_1 is one of $V(a)$ and let $V(d_1) = \{a, c, v_1\}$. If v_1 has degree 3, then $NV_3(a, c) + |V_4(a, c)| = 4$ and $V_Y(a, c) = \{v_1\}$, providing $|\widehat{E}_{a, c}| \leq 8$. Otherwise v_1 has degree 4 and it is different from d_2 and d_3 . For any $i = 2, 3$, each vertex of $V(d_i) \setminus \{c\}$ is either of degree 3 or v_1 . Thus $NE(d_2, d_3) = 8$ and $NV_3(d_2, d_3) + |V_4(d_2, d_3)| = 4$, providing $|\widehat{E}_{d_2, d_3}| \leq 9$. But $\widehat{G}_{d_2, d_3}$ has a triangle containing vertices a, b and d_1 . See Figure 6(a).

Last, assume that $|V_3(c)| \geq 2$ and let d_1 and d_2 be such vertices. As in the previous case, we may say that d_1 and d_2 are in $V(a)$, and $V(d_i) = \{a, c, v_i\}$ for $i = 1, 2$ where v_i has degree 4. When $v_1 = v_2$, $|V_3(a)| = 3$, $|V_4(a, c)| = 1$, and furthermore v_1 has degree 2 when we construct $\widehat{G}_{a, c}$, providing $|\widehat{E}_{a, c}| \leq 8$. When $\text{dist}(v_1, v_2) \geq 2$, three cases occur as follows; $|V_3(v_1)| \geq 3$, $|V_3(v_2)| \geq 3$, or for both $i = 1, 2$ $|V_3(v_i)| = 2$ and $V_4(v_i) = V_4 \setminus \{a, c, v_1, v_2\}$. All three cases satisfy that $NV_3(v_1, v_2) + |V_4(v_1, v_2)| \geq 4$, providing $|\widehat{E}_{v_1, v_2}| \leq 9$. But $\widehat{G}_{v_1, v_2}$ has a bigon containing vertices a and c . Finally when $\text{dist}(v_1, v_2) = 1$, two cases occur as follows. If d_3 has degree 3, then by the same reason as before we may say that d_3 is also in $V(a)$, and $V(d_3) = \{a, c, v_3\}$ where v_3 has degree 4. By the previous argument any pair of v_1, v_2 and v_3 has distance 1. This implies that G contains a triangle. If d_3 has degree 4, then $|V_3(d_3)| \geq 2$ because at most one vertex of $V(d_3)$ can be v_1 or v_2 . Thus $NV_3(a, d_3) \geq 4$, providing $|\widehat{E}_{a, d_3}| \leq 9$. But $\widehat{G}_{a, d_3}$ has a triangle containing vertices c, v_1 and v_2 . See Figure 6(b). $\square$




FIGURE 6. Some non-intrinsically knotted cases

Claim 2. If H is not a cycle with 6 edges, then G is not intrinsically knotted.

Proof. By Claim 1, if H is not a cycle with 6 edges, then H contains a cycle with 4 or 5 edges. First assume that H contains a cycle with 5 edges. Let $\{a_1, \dots, a_5\}$ be the set of five vertices of the cycle appeared in clockwise order. If the remaining vertex b of V_4 is contained in some $V(a_i)$, say $i = 1$, then b must have distance 1 from one of a_3 and a_4 , say a_3 , by Claim 1. See Figure 7(a). If $V_3(a_2) \neq V_3(b)$, $NV_3(a_2, b) + |V_4(a_2, b)| \geq 5$, providing $|\widehat{E}_{a_2, b}| \leq 8$. Otherwise $V_3(a_2) = V_3(b)$. Let c_1 and c_3 be the vertices of $V_3(a_1)$ and $V_3(a_3)$, respectively. If $c_1 = c_3$, we still have $|\widehat{E}_{a_2, b}| \leq 9$ and $\widehat{G}_{a_2, b}$ has a triangle containing vertices a_5 , a_4 and $c_1 = c_3$. If $c_1 \neq c_3$, then $|\widehat{E}_{a_1, a_3}| \leq 9$ and $\widehat{G}_{a_1, a_3}$ has a bigon as in the figure.

If b is not contained in $V(a_i)$ for any $i = 1, \dots, 5$, then $|V_3(a_i)| = 2$. If there is a pair of vertices a_i and a_{i+2} (or a_{i-3} if $i = 4, 5$) such that $V_3(a_i)$ and $V_3(a_{i+2})$ are disjoint, then $NV_3(a_i, a_{i+2}) + |V_4(a_i, a_{i+2})| = 5$, providing $|\widehat{E}_{a_i, a_{i+2}}| \leq 8$. Otherwise, for any pair of vertices a_i and a_{i+2} (or a_{i-3} if $i = 4, 5$), $V_3(a_i)$ and $V_3(a_{i+2})$ share vertices. Then they must share only one vertex as in Figure 7(b). Since there is only one extra vertex b of degree 4, for some pair of vertices a_i and a_{i+2} , $NV_3(a_i, a_{i+2}) + |V_4(a_i, a_{i+2})| = 4$ and $V_Y(a_i, a_{i+2}) \geq 1$, providing $|\widehat{E}_{a_i, a_{i+2}}| \leq 8$.




FIGURE 7. Cycle with 5 edges

Now assume that H contains a cycle with 4 edges. Let $\{a_1, \dots, a_4\}$ be the set of four vertices of the cycle appeared in clockwise order. If $V(a_1)$ and $V(a_3)$ (or similarly for $V(a_2)$ and $V(a_4)$) share only two vertices a_2 and a_4 , then the remaining two vertices of V_4 must be contained in $V(a_1) \cup V(a_3)$, otherwise $NV_3(a_1, a_3) \geq 3$ and $|V_4(a_1, a_3)| = 2$, providing $|\widehat{E}_{a_1, a_3}| \leq 8$. By Claim 1, the two vertices have distance 1, so H contains a cycle with 5 edges which was dealt in the previous case. If $V(a_1)$ and $V(a_3)$ (or similarly for $V(a_2)$ and $V(a_4)$) share exactly three vertices a_2 , a_4 and b , says, then let c_1 and c_3 be the remaining vertices of $V(a_1)$ and $V(a_3)$, respectively. If both c_1 and c_3 have degree 3, then $NV_3(a_1, a_3) + |V_4(a_1, a_3)| \geq 5$. If both have degree 4, then H contains a cycle with 5 edges as in the previous case. Finally if only c_1 (or similarly c_3) has degree 4, then, by Claim 1, $V(c_1)$ contains another vertex, say d , of V_4 and also d must have distance 1 from one of a_2 and a_4 , say a_4 , as in Figure 8(a). So $NV_3(a_4, c_1) + |V_4(a_4, c_1)| \geq 4$, providing $|\widehat{E}_{a_4, c_1}| \leq 9$, and $\widehat{G}_{a_4, c_1}$ has a triangle containing vertices a_2 , a_3 , and b . Now we may assume that $V(a_1) = V(a_3)$ and $V(a_2) = V(a_4)$. Then $NV_3(a_1, a_3) + |V_4(a_1, a_3)| = 4$, providing $|\widehat{E}_{a_1, a_3}| \leq 9$, and $\widehat{G}_{a_1, a_3}$ has a bigon as in Figure 8(b). $\square$

By Claim 2, H is exactly a cycle with 6 edges. Let $\{a_1, \dots, a_6\}$ be the set of six vertices of the cycle appeared in clockwise order. First suppose that there is not




FIGURE 8. Cycle with 4 edges


a vertex b in V_3 such that $V(b) = \{a_1, a_3, a_5\}$. If $V_3(a_1)$ and $V_3(a_3)$ are disjoint, then $NV_3(a_1, a_3) + |V_4(a_1, a_3)| = 5$. If $V_3(a_1)$ and $V_3(a_3)$ share exactly one vertex c , then the vertex of $V(c) \setminus \{a_1, a_3\}$ is not a_5 , so it should be one of $V_Y(a_1, a_3)$. Thus $NV_3(a_1, a_3) + |V_4(a_1, a_3)| + |V_Y(a_1, a_3)| = 5$. If $V_3(a_1)$ and $V_3(a_3)$ are same, then $NV_3(a_1, a_5) + |V_4(a_1, a_5)| = 5$ because $V_3(a_1)$ and $V_3(a_5)$ are disjoint. All three cases guarantee that G is not intrinsically knotted. Therefore we may assume that there are two vertices b_1 and b_2 so that $V(b_1) = \{a_1, a_3, a_5\}$ and $V(b_2) = \{a_2, a_4, a_6\}$. See Figure 9(a).

Suppose that there is a vertex c except b_1 so that $V(c)$ contains a_1 and a_3 . Let d_2 and d_5 be the vertices of $V_3(a_2)$ and $V_3(a_5)$, other than b_1 and b_2 , respectively. If $d_2 \neq d_5$, then $NV_3(a_2, a_5) = 4$. If $d_2 = d_5$, then $NV_3(a_2, a_5) = 3$ and $V_Y(a_2, a_5)$ is not empty. Both cases provide $|\hat{E}_{a_2, a_5}| \leq 9$, and $\hat{G}_{a_2, a_5}$ has a triangle containing vertices a_1, a_3 and c . Therefore we may assume in general that for any vertex c , except b_1 and b_2 , $V(c)$ does not contain both a_i and a_{i+2} for any $i = 1, 2, 3, 4$, and both a_i and a_{i-4} for any $i = 5, 6$.

Now consider $E \setminus \{E^2(b_1) \cup E^2(b_2)\}$ which consists of three extra edges. Note that each vertex of these edges has degree 3 and there are four more vertices of degree 3, besides b_1 and b_2 . These two facts guarantee that these extra edges must be connected as a tree. There are two types of the tree so as all three edges are incident to one vertex d or two edges are incident to different endpoints of the other edge e , respectively. In both cases, any two edges adjoined to the tree at the same vertex at the end must be also incident to a_i and a_{i+3} , respectively, for some $i = 1, 2, 3$. Therefore G is one of three graphs as in Figure 9(b)-(c), depending on the types of the tree. The graph G in Figure 9(b) is H_{12} which is intrinsically knotted. But the two graphs in Figure 9(c) are not intrinsically knotted because, for some i , $|\hat{E}_{a_i, a_{i+2}}| \leq 9$, and $\hat{G}_{a_i, a_{i+2}}$ has a triangle.

4.3. Case $(|V_4|, |V_3|) = (9, 2)$.

Let b_1 and b_2 be the vertices of V_3 . Since $|V_3| = 2$, there are at least three vertices a_1, a_2 and a_3 in V_4 such that all vertices of each $V(a_i)$ have degree 4. If $\text{dist}(a_1, a_2) = 1$, then $V(a_1) \cup V(a_2)$ consists of 8 vertices of V_4 , and so let c be the ninth vertex. Let d be any vertex among $V(a_1) \cup V(a_2) \setminus \{a_1, a_2\}$ which is not contained in $V(c)$. We assume that d is in $V(a_1)$. Then $V(d)$ should be contained in $V(a_2) \cup \{b_1, b_2\}$. This implies that $NE(a_2, d) = 8$ and $|V_3(d)| + |V_4(a_2, d)| \geq 4$, providing $|\hat{E}_{a_2, d}| \leq 9$. Since c has degree 4 in $\hat{G}_{a_2, d}$, it is not homeomorphic to $K(3, 3)$. We have the same result for any choices of pairs among a_1, a_2 , and a_3 .


FIGURE 9. Constructing H_{12}

Now assume that the distance between any pair among a_1, a_2 , and a_3 is at least 2. We separate into several cases according to the number $|V_4(a_1, a_2)|$. If $V_4(a_1, a_2) = \emptyset$ (i.e. $\text{dist}(a_1, a_2) > 2$), then $|V_4| \geq 10$, a contradiction. If $V_4(a_1, a_2) = \{d\}$, then $V_4 = V(a_1) \cup V(a_2) \cup \{a_1, a_2\}$. This implies that $a_3 \in V(a_1) \cup V(a_2)$, so $\text{dist}(a_1, a_3) = 1$ or $\text{dist}(a_2, a_3) = 1$ both of which was dealt in the previous case. If $V_4(a_1, a_2) = \{d_1, d_2\}$, then $V(d_1) \cup V(d_2) \setminus \{a_1, a_2\}$ is contained in $\{a_3, b_1, b_2\}$. This implies that each $V(d_i) \setminus \{a_1, a_2\}$ is the set of two vertices among $\{a_3, b_1, b_2\}$ so that $|V_3(d_1, d_2)| + |V_4(d_1, d_2)| \geq 4$, providing $|\hat{E}_{d_1, d_2}| \leq 9$. Since at least two of four vertices $V(a_1) \cup V(a_2) \setminus \{d_1, d_2\}$ still have degree 4 in $\hat{G}_{d_1, d_2}$, it is not homeomorphic to $K(3, 3)$. If $V_4(a_1, a_2) = \{d_1, d_2, d_3\}$, then $V(d_1) \cup V(d_2) \cup V(d_3) \setminus \{a_1, a_2\}$ is contained in $\{a_3, a_4, b_1, b_2\}$ where a_3 and a_4 are the remaining two vertices of degree 4 other than $V(a_1) \cup V(a_2) \cup \{a_1, a_2\}$. Thus each $V(d_i) \setminus \{a_1, a_2\}$ is the set of two vertices among $\{a_3, a_4, b_1, b_2\}$. This implies that $|V_3(d_i, d_j)| + |V_4(d_i, d_j)| \geq 4$ for some $i, j = 1, 2, 3$, providing $|\hat{E}_{d_i, d_j}| \leq 9$. Since at least one of three vertices $V(a_1) \cup V(a_2) \setminus \{d_i, d_j\}$ still has degree 4 in $\hat{G}_{d_i, d_j}$, it is not homeomorphic to $K(3, 3)$. Finally if $|V_4(a_1, a_2)| = 4$, then $|\hat{E}_{a_1, a_2}| \leq 9$. Since $\hat{G}_{a_1, a_2}$ still has the remaining three vertices of degree 4, it is not homeomorphic to $K(3, 3)$.

5. $\deg(a) = 3$

Since we are working on the graph with 21 edges and every vertex has degree 3, there is exactly 14 vertices. First suppose that there exists a pair of vertices a and b with $\text{dist}(a, b) \geq 4$. Then $E^2(a)$ and $E^2(b)$ can share only vertices but do not share edges in common. Since $|E^2(a) \cup E^2(b)| = 18$ and $|V(a) \cup V(b) \cup \{a, b\}| = 8$, the 18 endpoints of $E^2(a)$, $E^2(b)$ and three extra edges which are $E \setminus \{E^2(a) \cup E^2(b)\}$ meet at six vertices. If any two edges of $E^2(a) \setminus E(a)$ (and similarly for b) are incident to one vertex c of these six vertices, take the unique vertex d of $V(a)$ which is not the endpoints of these two edges. Then $NE(b, d) = 6$ and $NV_3(b, d) = 6$, providing

$|\widehat{E}_{b,d}| = 9$. But $\widehat{G}_{b,d}$ has a triangle containing c and the two vertices of $V(a) \setminus \{d\}$, so it is not homeomorphic to $K(3, 3)$. If not, each of these six vertices is common endpoints of one edge of $E^2(a)$, one edge of $E^2(b)$, and one extra edge. Now take an extra edge e and let b_1 and b_2 be the two vertices of $V(b)$ which have distance 1 from the endpoints of e . Let b_3 be the remaining vertex of $V(b)$. Then $NE(b_1, b_2) = 6$, $NV_3(b_1, b_2) = 5$, and $V_Y(b_1, b_2) = \{b_3\}$, providing $|\widehat{E}_{b_1, b_2}| = 9$. But $\widehat{G}_{b_1, b_2}$ has a triangle containing a and two vertices of $V(a)$, so it is not homeomorphic to $K(3, 3)$. See Figure 10(a).

FIGURE 10. Constructing C_{14}

Therefore we assume that the distance between any pair of vertices cannot exceed 3. Now we construct the intrinsically knotted graph G satisfying such conditions. Take a vertex a and let $V(a) = \{b_1, b_2, b_3\}$ and $V(b_i) = \{a, c_{2i-1}, c_{2i}\}$ for $i = 1, 2, 3$. As in Figure 10(b) the graph $E(a) \cup E(c_1) \cup \dots \cup E(c_6)$ consists of 21 edges and 22 vertices. This is the only way to draw the graph with 21 edges such that all vertices have distance at most 3 from a and 10 vertices $a, b_1, b_2, b_3, c_1, \dots, c_5$, and c_6 have degree 3. Now we join 12 vertices other than previous 10 vertices into 4 groups indicating the remaining 4 vertices d_1, d_2, d_3 and d_4 . Thus each $V(d_j)$, $j = 1, 2, 3, 4$, consists of three vertices among $c_1, \dots, c_6$. Since the distance between any c_i and $c_{i'}$ cannot exceed 3, the following two properties must be satisfied. First property is that $V(d_j)$ contains exactly one vertex from each group $\{c_{2i-1}, c_{2i}\}$ for $i = 1, 2, 3$. For example, if $V(d_1) = \{c_1, c_2, c_3\}$ (i.e. two vertices from the group $\{c_1, c_2\}$), then we can connect c_1 to at most two vertices among $\{c_4, c_5, c_6\}$ through some $E(d_j)$. This means that the distance between c_1 and one among $\{c_4, c_5, c_6\}$ exceed 3. Second is that different $V(d_j)$ and $V(d_{j'})$ share at most one vertex. For example, if they share two vertices c_1 and c_3 , then $\text{dist}(c_1, c_4) = 4$. From these two properties, without loss of generality, we may say that $V(d_1) = \{c_1, c_3, c_5\}$, $V(d_2) = \{c_1, c_4, c_6\}$, $V(d_3) = \{c_2, c_3, c_6\}$, and $V(d_4) = \{c_2, c_4, c_5\}$ as drawn in Figure 10(b). This graph is exactly C_{14} .

REFERENCES

- [1] P. Blain, G. Bowlin, T. Fleming, J. Foisy, J. Hendricks, and J. LaCombe, *Some results on intrinsically knotted graphs*, J. Knot Theory Ramif. **16** (2007) 749–760.
- [2] J. Conway and C. McA. Gordon, *Knots and links in spatial graphs*, J. Graph Theory **7** (1985) 445–453.
- [3] E. Flapan and R. Naimi, *The $Y\nabla$ moves does not preserve intrinsic knottedness*, Osaka J. Math. **45** (2008) 107–111.
- [4] J. Foisy, *Intrinsically knotted graphs*, J. Graph Theory **39** (2002) 178–187.
- [5] N. Goldberg, T. Mattman, and R. Naimi, *Many, many more intrinsically knotted graphs*, Preprint arXiv:1109.1632v1

- [6] R. Hanaki, R. Nikkuni, K. Taniyama, and A. Yamazaki, *On intrinsically knotted or completely 3-linked graphs*, Pacific J. Math. **252** (2011) 407–425.
- [7] B. Johnson, M. Kidwell, and T. Michael, *Intrinsically knotted graphs have at least 21 edges*, J. Knot Theory Ramif. **19** (2010) 1423–1429.
- [8] T. Kohara and S. Suzuki, *Some remarks on knots and links in spatial graphs*, Knots 90 (Osaka, 1990) (1992) 435–445.
- [9] R. Motwani, A. Raghunathan, and H. Saran, *Constructive results from graph minors; linkless embeddings*, Proc. 29th Annual Symposium on Foundations of Computer Science, IEEE (1988) 398–409.
- [10] M. Ozawa and Y. Tsutsumi, *Primitive spatial graphs and graph minors*, Rev. Mat. Complut. **20** (2007) 391–406.
- [11] N. Robertson and P. Seymour, *Graph minors XX, Wagner’s conjecture*, J. Combin. Theory Ser. B **92** (2004) 325–357.
- [12] H. Sachs, *On spatial representations of finite graphs*, Colloq. Math. Soc. Janos Bolyai **37** (1984) 649–662.

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, ANAM-DONG, SUNGBUK-KU, SEOUL 136-701, KOREA

E-mail address: mjmlj@korea.ac.kr

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, ANAM-DONG, SUNGBUK-KU, SEOUL 136-701, KOREA

E-mail address: Kimhjun@korea.ac.kr

DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, 84 HEUKSEOK-RO, DONGJAK-GU, SEOUL 156-756, KOREA

E-mail address: hjwith@cau.ac.kr

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, ANAM-DONG, SUNGBUK-KU, SEOUL 136-701, KOREA

E-mail address: seungsang@korea.ac.kr